



## Some Properties for Certain General Integral Operator

Vasile Marius Macarie<sup>1,\*</sup>, Daniel Breaz<sup>2</sup>

<sup>1</sup> Department of Mathematics and Computer Sciences, University of Pitești, Pitești, România

<sup>2</sup> Department of Mathematics, "1 Decembrie 1918" University of Alba Iulia, Alba Iulia, România

---

**Abstract.** In this paper we consider some subclasses of the class of analytic functions defined in the open unit disk of the complex plane and we study some properties for an integral operator on these classes. Particular results are presented.

**2010 Mathematics Subject Classifications:** 30C45

**Key Words and Phrases:** integral operator, analytic function, convex function, starlike function

---

### 1. Introduction

Let  $\mathcal{A}$  denote the class of the functions  $f$  of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

which are analytic in the open unit disk  $U = \{z \in \mathbb{C} : |z| < 1\}$ . We also denote by  $S$  the subclass of  $\mathcal{A}$  consisting of functions which are univalent in  $U$ .

A function  $f \in \mathcal{A}$  is said to be *convex of order*  $\alpha$ ,  $0 \leq \alpha < 1$  if it satisfies the condition

$$\operatorname{Re} \left( \frac{zf''(z)}{f'(z)} + 1 \right) > \alpha, \quad (z \in U)$$

and we denote this class by  $K(\alpha)$ .

A function  $f \in \mathcal{A}$  is said to be *starlike of order*  $\alpha$ ,  $0 \leq \alpha \leq 1$  if it satisfies the condition

$$\operatorname{Re} \left( \frac{zf'(z)}{f(z)} \right) > \alpha, \quad (z \in U)$$

and denote this class by  $S^*(\alpha)$ .

---

\*Corresponding author.

Email addresses: macariem@yahoo.com (V. Macarie), dbreaz@uab.ro (D. Breaz)

Let  $\mathcal{N}(\rho)$  be the subclass of  $\mathcal{A}$  consisting of the functions  $f$  which satisfy the inequality

$$\operatorname{Re} \left( 1 + \frac{zf''(z)}{f'(z)} \right) < \rho, \quad \rho > 1, \quad (z \in U).$$

This class was studied by S. Owa and H.M. Srivastava in [5].

A. Mohammed et al. considered in [4]  $\mathcal{MT}(\mu, \beta)$  the subclass of  $\mathcal{A}$  consisting of the functions  $f$  which satisfy the inequality

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < \beta \left| \mu \frac{zf'(z)}{f(z)} + 1 \right|, \quad 0 < \beta \leq 1, \quad 0 \leq \mu < 1, \quad (z \in U).$$

Also, Frasin and Jahangiri introduced in [2] the family  $\mathcal{B}(\mu, \alpha)$ ,  $\mu \geq 0$ ,  $0 \leq \alpha \leq 1$ , consisting of the functions  $f$  which satisfy the condition

$$\left| f'(z) \left( \frac{z}{f(z)} \right)^\mu - 1 \right| < 1 - \alpha, \quad (z \in U).$$

This family is a comprehensive class of analytic functions that includes various classes of analytic functions. We have  $\mathcal{B}(1, \alpha) \equiv S^*(\alpha)$  and  $\mathcal{B}(0, \alpha) \equiv R(\alpha)$ .

Let  $\beta - S_p(\alpha)$  be the subclass of  $\mathcal{A}$  consisting of the functions  $f$  which satisfy the inequality

$$\operatorname{Re} \left( \frac{zf'(z)}{f(z)} - \alpha \right) \geq \beta \left| \frac{zf'(z)}{f(z)} - 1 \right|, \quad -1 \leq \alpha \leq 1, \quad \beta > 0, \quad (z \in U).$$

This class was studied by M. Darus in [1].

A function  $f$  is said to be in the class  $KD(\mu, \alpha)$  if it satisfies the inequality

$$\operatorname{Re} \left( \frac{zf''(z)}{f'(z)} + 1 \right) \geq \mu \left| \frac{zf''(z)}{f'(z)} \right| + \alpha, \quad \mu \geq 0, \quad 0 \leq \alpha < 1, \quad (z \in U).$$

This class was studied by S. Shams et al. in [6].

In the present paper we study some properties for the integral operator  $G_n$  defined by

$$G_n(z) = \int_0^z \prod_{i=1}^n (f_i(t))^{\gamma_i-1} (g'_i(t))^{\eta_i} dt \tag{1}$$

on the classes presented above.

In order to prove our main results we need the following lemma:

**Lemma 1** (General Schwarz Lemma [3]). *Let the function  $f$  be regular in the disk  $U_R = \{z \in \mathbb{C} : |z| < R\}$ , with  $|f(z)| < M$  for fixed  $M$ . If  $f$  has one zero with multiplicity order bigger than  $m$  for  $z = 0$ , then*

$$|f(z)| \leq \frac{M}{R^m} \cdot |z|^m \quad (z \in U_R).$$

The equality can hold only if

$$f(z) = e^{i\theta} \cdot \frac{M}{R^m} \cdot z^m,$$

where  $\theta$  is constant.

### 2. Main Results

**Theorem 1.** Let  $\gamma_i \in \mathbb{R}, \gamma_i > 1, \eta_i \in \mathbb{R}, \eta_i > 0$  for all  $i = 1, 2, \dots, n$ , the functions  $f_i \in \mathcal{MT}(\mu_i, \beta_i), 0 < \beta_i \leq 1, 0 \leq \mu_i < 1$  and  $g_i \in \mathcal{A}$  for all  $i = 1, 2, \dots, n$  satisfying the conditions

$$\left| \frac{f'_i(z)}{f_i(z)} \right| < M_i, (M_i \geq 1) \text{ for all } i = 1, 2, \dots, n \tag{2}$$

and

$$\left| \frac{g''_i(z)}{g'_i(z)} \right| < N_i, (N_i \geq 1) \text{ for all } i = 1, 2, \dots, n. \tag{3}$$

Then the integral operator  $G_n$  defined in (1) is in  $\mathcal{N}(\rho)$ , where

$$\rho = 1 + \sum_{i=1}^n [(\gamma_i - 1)(\beta_i \mu_i M_i + \beta_i + 1) + \eta_i N_i]$$

*Proof.* From (1), we have

$$G'_n(z) = \prod_{i=1}^n (f_i(z))^{\gamma_i - 1} (g'_i(z))^{\eta_i}$$

and

$$\frac{zG''_n(z)}{G'_n(z)} = \sum_{i=1}^n (\gamma_i - 1) \frac{zf'_i(z)}{f_i(z)} + \sum_{i=1}^n \eta_i \frac{zg''_i(z)}{g'_i(z)}.$$

Thus, we have

$$\operatorname{Re} \left( \frac{zG''_n(z)}{G'_n(z)} + 1 \right) = \sum_{i=1}^n (\gamma_i - 1) \operatorname{Re} \left( \frac{zf'_i(z)}{f_i(z)} \right) + \sum_{i=1}^n \eta_i \operatorname{Re} \left( \frac{zg''_i(z)}{g'_i(z)} \right) + 1.$$

Since  $\operatorname{Re} w \leq |w|$ , then

$$\operatorname{Re} \left( \frac{zG''_n(z)}{G'_n(z)} + 1 \right) \leq \sum_{i=1}^n (\gamma_i - 1) \left| \frac{zf'_i(z)}{f_i(z)} \right| + \sum_{i=1}^n \eta_i \left| \frac{zg''_i(z)}{g'_i(z)} \right| + 1. \tag{4}$$

Using that  $f_i \in \mathcal{MT}(\mu_i, \beta_i)$  for all  $i = 1, 2, \dots, n$  in relation (4), we obtain

$$\begin{aligned} \operatorname{Re} \left( \frac{zG''_n(z)}{G'_n(z)} + 1 \right) &\leq \sum_{i=1}^n (\gamma_i - 1) \left( \left| \frac{zf'_i(z)}{f_i(z)} - 1 \right| + 1 \right) + \sum_{i=1}^n \eta_i \left| \frac{zg''_i(z)}{g'_i(z)} \right| + 1 \\ &< \sum_{i=1}^n (\gamma_i - 1) \beta_i \left| \mu_i \frac{zf'_i(z)}{f_i(z)} + 1 \right| + \sum_{i=1}^n (\gamma_i - 1) \\ &\quad + \sum_{i=1}^n \eta_i \left| \frac{zg''_i(z)}{g'_i(z)} \right| + 1 \end{aligned}$$

and using the hypothesis (2) and (3) in this last relation we have

$$\operatorname{Re} \left( \frac{zG_n''(z)}{G_n'(z)} + 1 \right) < \sum_{i=1}^n [(\gamma_i - 1)(\beta_i \mu_i M_i + \beta_i + 1) + \eta_i N_i] + 1 = \rho$$

This completes the proof of our theorem. □

Letting  $n = 1$ ,  $\gamma_1 = \gamma$ ,  $\eta_1 = \eta$ ,  $M_1 = M$ ,  $N_1 = N$ ,  $\mu_1 = \mu$ ,  $\beta_1 = \beta$ ,  $f_1 = f$  and  $g_1 = g$  in Theorem 1, we have

**Corollary 1.** Let  $\gamma \in \mathbb{R}$ ,  $\gamma > 1$ ,  $\eta \in \mathbb{R}$ ,  $\eta > 0$ , the functions  $f \in \mathcal{MT}(\mu, \beta)$ ,  $0 < \beta \leq 1$ ,  $0 \leq \mu < 1$  and  $g \in \mathcal{A}$  satisfying the conditions

$$\left| \frac{f'(z)}{f(z)} \right| < M, (M \geq 1) \text{ and } \left| \frac{g''(z)}{g'(z)} \right| < N, (N \geq 1).$$

Then the integral operator

$$G_1(z) = \int_0^z (f(t))^{\gamma-1} (g'(t))^\eta dt$$

is in  $\mathcal{N}(\rho)$ , where  $\rho = (\gamma - 1)(\beta\mu M + \beta + 1) + \eta N + 1$ .

Letting  $\gamma = 2$ ,  $\eta = 1$  in Corollary 1, we have

**Corollary 2.** Let  $f \in \mathcal{MT}(\mu, \beta)$ ,  $0 < \beta \leq 1$ ,  $0 \leq \mu < 1$  and  $g \in \mathcal{A}$  satisfying the conditions

$$\left| \frac{f'(z)}{f(z)} \right| < M, (M \geq 1) \text{ and } \left| \frac{g''(z)}{g'(z)} \right| < N, (N \geq 1).$$

Then the integral operator

$$G(z) = \int_0^z f(t)g'(t)dt$$

is in  $\mathcal{N}(\rho)$ , where  $\rho = \beta(\mu M + 1) + N + 2$ .

**Theorem 2.** Let  $\gamma_i \in \mathbb{R}$ ,  $\gamma_i > 1$ ,  $\eta_i \in \mathbb{R}$ ,  $\eta_i > 0$  for all  $i = 1, 2, \dots, n$ , the functions  $f_i \in \mathcal{B}(\mu_i, \alpha_i)$ ,  $\mu_i \geq 0$ ,  $0 \leq \alpha_i < 1$  satisfying the conditions  $|f_i(z)| \leq M_i$ ,  $(M_i \geq 1)$  and  $g_i \in \mathcal{N}(\rho_i)$ ,  $\rho_i > 1$  for all  $i = 1, 2, \dots, n$ . If

$$\sum_{i=1}^n [(\gamma_i - 1)(2 - \alpha_i)M_i^{\mu_i-1} + \eta_i(\rho_i - 1)] < 1,$$

then the integral operator  $G_n$  defined in (1) is in  $K(\delta)$ , where

$$\delta = 1 - \sum_{i=1}^n [(\gamma_i - 1)(2 - \alpha_i)M_i^{\mu_i-1} + \eta_i(\rho_i - 1)].$$

*Proof.* From (1), we have

$$G'_n(z) = \prod_{i=1}^n (f_i(z))^{\gamma_i-1} (g'_i(z))^{\eta_i}$$

and

$$\frac{zG''_n(z)}{G'_n(z)} = \sum_{i=1}^n (\gamma_i - 1) \frac{zf'_i(z)}{f_i(z)} + \sum_{i=1}^n \eta_i \frac{zg''_i(z)}{g'_i(z)}.$$

Hence

$$\begin{aligned} \left| \frac{zG''_n(z)}{G'_n(z)} \right| &\leq \sum_{i=1}^n (\gamma_i - 1) \left( \left| f'_i(z) \left( \frac{z}{f_i(z)} \right)^{\mu_i} - 1 \right| + 1 \right) \left| \frac{f_i(z)}{z} \right|^{\mu_i-1} \\ &\quad + \sum_{i=1}^n \eta_i \left( \left| \frac{zg''_i(z)}{g'_i(z)} + 1 \right| - 1 \right) \end{aligned} \tag{5}$$

Since  $|f_i(z)| \leq M_i$  for all  $i = 1, 2, \dots, n$ , applying the General Schwarz Lemma, it results

$$\left| \frac{f_i(z)}{z} \right| \leq M_i \text{ for all } i = 1, 2, \dots, n. \tag{6}$$

From (5) and (6), using that  $f_i \in \mathcal{B}(\mu_i, \alpha_i)$  and  $g_i \in \mathcal{N}(\rho_i)$  for all  $i = 1, 2, \dots, n$ , we obtain

$$\left| \frac{zG''_n(z)}{G'_n(z)} \right| < \sum_{i=1}^n [(\gamma_i - 1)(2 - \alpha_i)M_i^{\mu_i-1} + \eta_i(\rho_i - 1)] = 1 - \delta$$

This completes the proof of our theorem. □

Letting  $n = 1$ ,  $\gamma_1 = \gamma$ ,  $\eta_1 = \eta$ ,  $M_1 = M$ ,  $\mu_1 = \mu$ ,  $\alpha_1 = \alpha$ ,  $\rho_1 = \rho$ ,  $f_1 = f$  and  $g_1 = g$  in Theorem 2, we have

**Corollary 3.** *Let  $\gamma \in \mathbb{R}$ ,  $\gamma > 1$ ,  $\eta \in \mathbb{R}$ ,  $\eta > 0$ , the functions  $f \in \mathcal{B}(\mu, \alpha)$ ,  $\mu \geq 0$ ,  $0 \leq \alpha < 1$ , satisfying the condition  $|f(z)| \leq M$ , ( $M \geq 1$ ) and  $g \in \mathcal{N}(\rho)$ ,  $\rho > 1$ . If*

$$(\gamma - 1)(2 - \alpha)M^{\mu-1} + \eta(\rho - 1) < 1$$

*then the integral operator*

$$G_1(z) = \int_0^z (f(t))^{\gamma-1} (g'(t))^\eta dt$$

*is in  $K(\delta)$ , where  $\delta = 1 + (\gamma - 1)(\alpha - 2)M^{\mu-1} + \eta(1 - \rho)$ .*

Letting  $\mu_i = 0$  and  $M_i = M$  for all  $i = 1, 2, \dots, n$  in Theorem 2, we have

**Corollary 4.** Let  $\gamma_i \in \mathbb{R}, \gamma_i > 1, \eta_i \in \mathbb{R}, \eta_i > 0$  for all  $i = 1, 2, \dots, n$ , the functions  $f_i \in R(\alpha_i), 0 \leq \alpha_i < 1$ , satisfying the conditions  $|f_i(z)| \leq M, (M \geq 1)$  and  $g_i \in \mathcal{N}(\rho_i), \rho_i > 1$  for all  $i = 1, 2, \dots, n$ . If

$$\sum_{i=1}^n \left[ (\gamma_i - 1)(2 - \alpha_i) \frac{1}{M} + \eta_i(\rho_i - 1) \right] < 1$$

then the integral operator  $G_n$  defined in (1) is in  $K(\delta)$ , where

$$\delta = 1 - \sum_{i=1}^n \left[ (\gamma_i - 1)(2 - \alpha_i) \frac{1}{M} + \eta_i(\rho_i - 1) \right].$$

Letting  $\mu_i = 1$  and  $M_i = M$  for all  $i = 1, 2, \dots, n$  in Theorem 2, we have

**Corollary 5.** Let  $\gamma_i \in \mathbb{R}, \gamma_i > 1, \eta_i \in \mathbb{R}, \eta_i > 0$  for all  $i = 1, 2, \dots, n$ , the functions  $f_i \in S^*(\alpha_i), 0 \leq \alpha_i < 1$ , satisfying the conditions  $|f_i(z)| \leq M, (M \geq 1)$  and  $g_i \in \mathcal{N}(\rho_i), \rho_i > 1$  for all  $i = 1, 2, \dots, n$ . If

$$\sum_{i=1}^n [(\gamma_i - 1)(2 - \alpha_i) + \eta_i(\rho_i - 1)] < 1$$

then the integral operator  $G_n$  defined in (1) is in  $K(\delta)$ , where

$$\delta = 1 - \sum_{i=1}^n [(\gamma_i - 1)(2 - \alpha_i) + \eta_i(\rho_i - 1)].$$

**Theorem 3.** Let  $\gamma_i \in \mathbb{R}, \gamma_i > 1, \eta_i \in \mathbb{R}, \eta_i > 0$  for all  $i = 1, 2, \dots, n$ , the functions  $f_i \in \rho_i - S_p(\varepsilon_i), -1 \leq \varepsilon_i \leq 1, \rho_i > 0$  and  $g_i \in KD(\mu_i, \alpha_i), 0 \leq \alpha_i < 1, \mu_i \geq 0$  for all  $i = 1, 2, \dots, n$ . If

$$0 < \sum_{i=1}^n [(1 - \gamma_i)\varepsilon_i + \eta_i(1 - \alpha_i)] \leq 1$$

then the integral operator  $G_n$  defined in (1) is in  $K(\delta)$ , where

$$\delta = 1 + \sum_{i=1}^n [(\gamma_i - 1)\varepsilon_i + \eta_i(\alpha_i - 1)].$$

*Proof.* Following the same steps as in Theorem 1, we obtain that

$$\frac{zG_n''(z)}{G_n'(z)} = \sum_{i=1}^n (\gamma_i - 1) \frac{zf_i'(z)}{f_i(z)} + \sum_{i=1}^n \eta_i \frac{zg_i''(z)}{g_i'(z)}$$

and hence

$$\begin{aligned} \frac{zG_n''(z)}{G_n'(z)} + 1 &= \sum_{i=1}^n \left[ (\gamma_i - 1) \left( \frac{zf_i'(z)}{f_i(z)} - \varepsilon_i \right) + (\gamma_i - 1)\varepsilon_i \right] \\ &\quad + \sum_{i=1}^n \left[ \eta_i \left( \frac{zg_i''(z)}{g_i'(z)} + 1 \right) - \eta_i \right] + 1. \end{aligned}$$

We calculate the real part from both terms of the above expression and obtain

$$\begin{aligned} \operatorname{Re} \left( \frac{zG_n''(z)}{G_n'(z)} + 1 \right) &= \sum_{i=1}^n (\gamma_i - 1) \operatorname{Re} \left( \frac{zf_i'(z)}{f_i(z)} - \varepsilon_i \right) + \sum_{i=1}^n (\gamma_i - 1) \varepsilon_i \\ &\quad + \sum_{i=1}^n \eta_i \operatorname{Re} \left( \frac{zg_i''(z)}{g_i'(z)} + 1 \right) - \sum_{i=1}^n \eta_i + 1. \end{aligned} \tag{7}$$

From (7), using that  $f_i \in \rho_i - S_p(\varepsilon_i)$  and  $g_i \in KD(\mu_i, \alpha_i)$  for all  $i = 1, 2, \dots, n$ , we have

$$\begin{aligned} \operatorname{Re} \left( \frac{zG_n''(z)}{G_n'(z)} + 1 \right) &\geq \sum_{i=1}^n (\gamma_i - 1) \rho_i \left| \frac{zf_i'(z)}{f_i(z)} - 1 \right| + \sum_{i=1}^n (\gamma_i - 1) \varepsilon_i \\ &\quad + \sum_{i=1}^n \eta_i \left( \mu_i \left| \frac{zg_i''(z)}{g_i'(z)} \right| + \alpha_i \right) - \sum_{i=1}^n \eta_i + 1. \end{aligned}$$

Then

$$\begin{aligned} \operatorname{Re} \left( \frac{zG_n''(z)}{G_n'(z)} + 1 \right) &\geq \sum_{i=1}^n (\gamma_i - 1) \rho_i \left| \frac{zf_i'(z)}{f_i(z)} - 1 \right| + \sum_{i=1}^n (\gamma_i - 1) \varepsilon_i \\ &\quad + \sum_{i=1}^n \eta_i \mu_i \left| \frac{zg_i''(z)}{g_i'(z)} \right| + \sum_{i=1}^n \eta_i (\alpha_i - 1) + 1. \end{aligned} \tag{8}$$

Since  $(\gamma_i - 1) \rho_i \left| \frac{zf_i'(z)}{f_i(z)} - 1 \right| > 0$  and  $\eta_i \mu_i \left| \frac{zg_i''(z)}{g_i'(z)} \right| \geq 0$  for all  $i = 1, 2, \dots, n$ , we obtain from (8) that

$$\operatorname{Re} \left( \frac{zG_n''(z)}{G_n'(z)} + 1 \right) > \sum_{i=1}^n [(\gamma_i - 1) \varepsilon_i + \eta_i (\alpha_i - 1)] + 1 = \delta.$$

This completes the proof of our theorem. □

Letting  $n = 1$ ,  $\gamma_1 = \gamma$ ,  $\eta_1 = \eta$ ,  $\rho_1 = \rho$ ,  $\varepsilon_1 = \varepsilon$ ,  $\mu_1 = \mu$ ,  $\alpha_1 = \alpha$ ,  $f_1 = f$  and  $g_1 = g$  in Theorem 3, we have

**Corollary 6.** *Let  $\gamma \in \mathbb{R}$ ,  $\gamma > 1$ ,  $\eta \in \mathbb{R}$ ,  $\eta > 0$ , the functions  $f \in \rho - S_p(\varepsilon)$ ,  $-1 \leq \varepsilon \leq 1$ ,  $\rho > 0$  and  $g \in KD(\mu, \alpha)$ ,  $0 \leq \alpha < 1$ ,  $\mu \geq 0$ . If*

$$0 < (1 - \gamma)\varepsilon + \eta(1 - \alpha) \leq 1$$

then the integral operator

$$G_1(z) = \int_0^z (f(t))^{\gamma-1} (g'(t))^\eta dt$$

is in  $K(\delta)$ , where  $\delta = 1 + (\gamma - 1)\varepsilon + \eta(\alpha - 1)$ .

Letting  $\gamma = 2$  and  $\eta = 1$  in Corollary 6, we have

**Corollary 7.** Let the functions  $f \in \rho - S_p(\varepsilon)$ ,  $-1 \leq \varepsilon \leq 1$ ,  $\rho > 0$  and  $g \in KD(\mu, \alpha)$ ,  $0 \leq \alpha < 1$ ,  $\mu \geq 0$ . If

$$0 < 1 - \alpha - \varepsilon \leq 1$$

then the integral operator

$$G(z) = \int_0^z f(t)g'(t)dt$$

is in  $K(\delta)$ , where  $\delta = \varepsilon + \alpha$ .

### References

- [1] M Darus. Certain class of uniformly analytic functions. *Acta Mathematica Academiae Pedagogicae Nyregyhaziensis*, 24:354-358, 2008.
- [2] B A Frasin and J M Jahangiri. A new and comprehensive class of analytic functions. *Annals of Oradea University-Mathematics Fascicola*, XV:59-62, 2008.
- [3] O Mayer. *The functions theory of one variable complex*. Bucuresti, 1981.
- [4] A Mohammed, M Darus, and D Breaz, Some properties for certain integral operators. *Acta Universitatis Apulensis*, 23:79-89, 2010.
- [5] S Owa and H M Srivastava. Some generalized convolution properties associated with certain subclasses of analytic functions. *Journal of Inequalities in Pure and Applied Mathematics*, 3(3):42:1-13, 2003.
- [6] S Shams, S R Kulkarni, and J M Jahangiri. Classes of uniformly starlike and convex functions. *International Journal of Mathematics and Mathematical Sciences*, 55:2959-2961, 2004.